






## RESEARCH ARTICLE

# High levels of eicosanoids and docosanoids in the lungs of intubated COVID-19 patients

Anne-Sophie Archambault<sup>1,2</sup>  | Younes Zaid<sup>3,4</sup>  | Volatiana Rakotoarivelo<sup>1,2</sup> |  
 Caroline Turcotte<sup>1,2</sup> | Étienne Doré<sup>5,6</sup> | Isabelle Dubuc<sup>5</sup> | Cyril Martin<sup>1,2</sup> |  
 Olivier Flamand<sup>5</sup> | Youssef Amar<sup>7</sup> | Amine Cheikh<sup>4</sup> | Hakima Fares<sup>4</sup> |  
 Amine El Hassani<sup>4</sup> | Youssef Tijani<sup>8</sup> | Andréanne Côté<sup>1</sup> | Michel Laviolette<sup>1</sup> |  
 Éric Boilard<sup>5,6,9</sup>  | Louis Flamand<sup>5,9</sup>  | Nicolas Flamand<sup>1,2</sup> 

<sup>1</sup>Centre de Recherche de l'Institut universitaire de cardiologie et de pneumologie de Québec, Faculté de médecine, Département de médecine, Université Laval, Québec, QC, Canada

<sup>2</sup>Canada Excellence Research Chair in the Microbiome-Endocannabinoidome Axis in Metabolic Health, Université Laval, Québec, QC, Canada

<sup>3</sup>Biology Department, Faculty of Sciences, Mohammed V University, Rabat, Morocco

<sup>4</sup>Cheikh Zaïd Hospital, Abulcasis University of Health Sciences, Rabat, Morocco

<sup>5</sup>Centre de Recherche du Centre Hospitalier, Universitaire de Québec-Université Laval, Québec, QC, Canada

<sup>6</sup>Centre de Recherche Arthrite, Université Laval, Québec, QC, Canada

<sup>7</sup>Moroccan Foundation for Advanced Science, Innovation & Research (MAScIR), Rabat, Morocco

<sup>8</sup>Faculty of Medicine, Mohammed VI University of Health Sciences, Casablanca, Morocco

<sup>9</sup>Département de Microbiologie-Inféctiologie et d'immunologie, Université Laval, Québec, QC, Canada

## Correspondence

Nicolas Flamand, Centre de Recherche de l'Institut universitaire de cardiologie et de pneumologie de Québec, Faculté de médecine, Département de médecine, Université Laval, 2725 chemin Ste-Foy, Office A3140, Québec, QC G1V 4G5, Canada.  
 Email: Nicolas.Flamand@criucpq.ulaval.ca

Louis Flamand, Centre de Recherche du Centre Hospitalier Universitaire de Québec-Université Laval, 2705 Laurier boulevard, Office T1-64, Québec, QC G1V 4G2, Canada.  
 Email: Louis.Flamand@crchudequebec.ulaval.ca

## Abstract

Severe acute respiratory syndrome coronavirus 2 is responsible for coronavirus disease 2019 (COVID-19). While COVID-19 is often benign, a subset of patients develops severe multilobar pneumonia that can progress to an acute respiratory distress syndrome. There is no cure for severe COVID-19 and few treatments significantly improved clinical outcome. Dexamethasone and possibly aspirin, which directly/indirectly target the biosynthesis/effects of numerous lipid mediators are among those options. Our objective was to define if severe COVID-19 patients were characterized by increased bioactive lipids modulating lung inflammation. A targeted lipidomic analysis of bronchoalveolar lavages (BALs) by tandem mass spectrometry was done on 25 healthy controls and 33 COVID-19 patients requiring mechanical ventilation.

**Abbreviations:** AA, arachidonic acid; ALA, linolenic acid; BAL, bronchoalveolar lavage; DGLA, dihomo- $\gamma$ -linolenic acid; DHA, docosahexaenoic acid; DPA, docosapentaenoic acid; EPA, eicosapentaenoic acid; EX, eoxin; HDHA, hydroxy-docosahexaenoic acid; HETE, hydroxyeicosatetraenoic acid; HEPE, hydroxyeicosapentaenoic acid; 12-HHTrE, 12-hydroxy-heptadecatrienoic acid; HODE, hydroxyoctadecadienoic acid; KETE, oxo-eicosatetraenoic acid; KODE, oxo-octadecadienoic acid; LA, linoleic acid; LO, lipoxygenase; LT, leukotriene; LX, lipoxin; PDX, protectin D1; PG, prostaglandin; Rv, resolvin; SPM, specialized pro-resolving mediators; TX, thromboxane.

Anne-Sophie Archambault and Younes Zaid are co-first authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *The FASEB Journal* published by Wiley Periodicals LLC on behalf of Federation of American Societies for Experimental Biology.

**Funding information**

Cheikh Zaid Foundation; Canadian New Frontier Research Fund, Grant/Award Number: NFRN-2019-00004; Natural Sciences and Engineering Research Council of Canada; Canadian Institutes of Health Research; Fonds de Recherche du Québec en Santé (FRQS)

BALs from severe COVID-19 patients were characterized by increased fatty acids and inflammatory lipid mediators. There was a predominance of thromboxane and prostaglandins. Leukotrienes were also increased, notably LTB<sub>4</sub>, LTE<sub>4</sub>, and eoxin E<sub>4</sub>. Monohydroxylated 15-lipoxygenase metabolites derived from linoleate, arachidonate, eicosapentaenoate, and docosahexaenoate were also increased. Finally yet importantly, specialized pro-resolving mediators, notably lipoxin A<sub>4</sub> and the D-series resolvins, were also increased, underscoring that the lipid mediator storm occurring in severe COVID-19 involves pro- and anti-inflammatory lipids. Our data unmask the lipid mediator storm occurring in the lungs of patients afflicted with severe COVID-19. We discuss which clinically available drugs could be helpful at modulating the lipidome we observed in the hope of minimizing the deleterious effects of pro-inflammatory lipids and enhancing the effects of anti-inflammatory and/or pro-resolving lipid mediators.

**KEYWORDS**

COVID-19, docosanoids, eicosanoids, eoxins, specialized pro-resolving mediators, thromboxane

## 1 | INTRODUCTION

Coronavirus disease 2019 (COVID-19) is the infectious disease caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2).<sup>1</sup> SARS-CoV-2 spreads predominantly from respiratory droplets of infected individuals to epithelial cells in the upper airways and oral cavity.<sup>2</sup> COVID-19 manifestations are broad, from asymptomatic carriers to severe viral pneumonia leading to respiratory failure and ultimately acute respiratory distress syndrome (ARDS) requiring mechanical ventilation.<sup>3,4</sup> In some patients, there is a yet to be explained hyperinflammatory response.

Autopsies from severe COVID-19 patients unmasked a heterogeneous disease that can be characterized by diffuse alveolar and/or epithelial damage, thrombosis, pulmonary embolism, and lymphocytic and/or granulocytic inflammatory cell infiltration.<sup>4-6</sup> An uncontrolled systemic inflammatory response known as the cytokine storm is suggested as an important contributor of SARS-CoV-2 lethality and is the consequence of an important release of cytokines such as tumor necrosis factor- $\alpha$ , interleukin (IL)-1 $\beta$ , and IL-6,<sup>4,7</sup> although this remains unclear.<sup>8</sup> The disease heterogeneity suggests that, aside cytokines/chemokines, other immunological effectors contribute to the sustained inflammatory responses observed in severe COVID-19 cases.

Bioactive lipids, notably eicosanoids, likely promote their fair share of deleterious effects by enhancing leukocyte recruitment/activation, promoting the exudate formation and by stimulating platelet aggregation and thrombus formation. In contrast, specialized pro-resolving mediators (SPMs),

mainly consisting of docosanoids, could dampen inflammation and promote its resolution.<sup>9</sup> Of note, some of these lipids are elevated in the blood of COVID-19 patients.<sup>10,11</sup> While therapeutic approaches targeting the biosynthesis/effects of inflammatory lipids are readily available, the levels of those lipids in the lungs during severe COVID-19 were unknown.

Herein, we performed a targeted lipidomic analysis of bronchoalveolar lavages fluids (BALs) from healthy volunteers and severe COVID-19 patients. Numerous eicosanoids, notably thromboxane, and SPMs were found in severe COVID-19 patients' BALs. Our study further highlights that few associations exist between clinical parameters and the bioactive lipids present in the primary site of disease, the lungs.

## 2 | METHODS

### 2.1 | Materials

All lipids were obtained from Cayman Chemical (Ann Arbor, MI, USA) and all LC-MS grade solvents were purchased from Fisher Scientific (Ottawa, Canada).

### 2.2 | Ethics

This study was approved by the local ethics committee (Cheikh Zaid Hospital in Rabat Morocco (Project: CEFCZ/PR/2020- PR04); Québec City Heart & Lung Institute; CHU de Québec-Université Laval). The study complies with the

Declaration of Helsinki and all subjects signed a consent form.

### 2.3 | Subject selection

Non-smoking healthy subjects taking no other medication than anovulants, without any documented acute or chronic inflammatory disease and without recent (8 weeks) airway infection were recruited for bronchoalveolar lavages at the Québec City Heart & Lung Institute (Canada). Severe COVID-19 patients were enrolled based on the inclusion criteria for intubation and the need of mechanical ventilation at Cheikh Zaid Hospital (Rabat, Morocco) between May and June 2020, before COVID-19 therapies such as remdesivir and dexamethasone were greenlit for treating COVID-19. Within 2 hours after intubation and the initiation of mechanical ventilation, blood sampling was done to assess the clinical variables (Table 1, Table S1 for each individual subjects). Patients were ventilated in the prone decubitus position with a tidal volume of 6 mL/kg and a PEEP varying between 8 and 14 cm H<sub>2</sub>O. The FiO<sub>2</sub> was adjusted between 0.6 and 1.0 to obtain a SpO<sub>2</sub> ≥ 92%.

### 2.4 | Bronchoalveolar lavages

BAL fluids from healthy volunteers were obtained as follows: One 50 mL bolus of sterile 0.9% saline was injected in a sub-segmental bronchi of the right middle lobe. The obtained lavages were centrifuged (350 g, 10 minutes) to pellet cells and the supernatants were immediately frozen (−80° Celsius) until further processing. A 5 mL aliquot of the samples was thawed then reduced to ~500 µL using a stream of nitrogen. BAL fluids from severe COVID-19 patients were obtained less than 2 hours following intubation as follows: a total of 100 mL of sterile 0.9% saline (2 boli of 50 mL) was injected in a sub-segmental bronchi of the right middle lobe. The obtained lavages were pooled and centrifuged (320 g, 15 minutes) to pellet cells. Supernatants were next concentrated as for the BALs of healthy donors and immediately frozen (−80° Celsius) until further processing.

### 2.5 | LC-MS/MS analyses

Concentrated samples (500 µL) were denatured with 500 µL of LC-MS-grade MeOH containing the internal standards then warmed at 60° Celsius for 30 minutes to inactivate SARS-CoV-2, denatured overnight (−20° Celsius), and then centrifuged (10,000 g) to remove the denatured proteins. Under these conditions, no significant degradation of

**TABLE 1** Clinical characteristics of BAL donors

	COVID-19	Healthy controls
N	33	25
Women/Men	16/17	16/9
Smoker (Y/N)	5/28	0/25
Death (Y/N)	2/31	N/A
SARS-CoV-2 detection in BAL (Ct)	25 ± 1	N/A
Age	58 ± 3	26 ± 1
Weight (kg)	73 ± 3	N/A
Hospital stay (days)	21 ± 2	N/A
Days before BAL	3 ± 1	N/A
D-Dimers (mg/mL)	1.21 ± 0.13	N/A
CRP (mg/l)	20.65 ± 2.53	N/A
ALT (U/l)	32.6 ± 2.1	N/A
AST (U/l)	35.1 ± 3.0	N/A
LDH (U/l)	616 ± 41	N/A
Hemoglobin (g/l)	123 ± 2	N/A
Blood Monocytes (million/mL)	0.3 ± 0.03	N/A
Blood Neutrophils (million/mL)	2.9 ± 0.19	N/A
Blood Eosinophils (million/mL)	0.03 ± 0.003	N/A
Blood Platelets (million/mL)	182 ± 13	N/A
Blood Lymphocytes (million/mL)	1.13 ± 0.13	N/A
Platelet to Lymphocyte ratio (PLR)	223.89 ± 34.93	N/A
BAL Neutrophils (million/mL)	24.7 ± 2.2	0.001 ± 0.0002
BAL Eosinophils (million/mL)	0.2 ± 0.08	0.000 ± 0.000
BAL Lymphocytes (million/mL)	21.8 ± 2.4	0.004 ± 0.001
BAL macrophages (million/mL)	N/A	0.073 ± 0.007

Note: Data are presented as the mean ± SEM.

Abbreviations: ALT, alanine transaminase; AST, aspartate transaminase; CRP, C-reactive protein; LDH, Lactate dehydrogenase; N/A, not available.

internal standards and selected lipids and lipid mediators was observed (Figure S1). Supernatants were diluted with water containing 0.01% acetic acid and lipids were extracted by solid phase extraction using Strata-X cartridges (Polymeric Reversed Phase, Phenomenex, USA) as described before.<sup>12</sup> Lipids were separated using the same column and LC program as in,<sup>12</sup> quantification was done using a Shimadzu 8050 triple quadrupole mass spectrometer and analyses were performed using multiple reaction monitoring for the specific mass transitions of each lipid as well as the deuterated

internal standard used (Table S2). Due to the large number of analytes being quantitated and/or commercial availability, some lipid mediators were analyzed with a surrogate, deuterated internal standard.


## 2.6 | Statistical analysis

Statistical analysis was made using the GraphPad Prism 9 software. The normality assumption was verified using the Shapiro-Wilk test and p-values were calculated using the Mann-Whitney test. Spearman correlations were calculated using *corr.test* function from *pshych* CRAN package and the correlation matrices were generated using the *corrplot* package. Values of  $P < .05$  were considered significant.

## 3 | RESULTS

While the cytokine profile observed in severe COVID-19 patients is relatively well-defined,<sup>4,13-16</sup> no study investigated the lipid content of BALs. We thus performed a targeted lipidomic analysis to define whether lipid mediators were present/increased in the lungs of severe COVID-19 patients.

In contrast to the BALs of healthy controls, numerous eicosanoids were detected in those of severe COVID-19 patients. As such, BALs from COVID-19 patients had increased AA levels as well as all measured COX metabolites (Figure 1B-D). The most increased COX metabolites were  $TXB_2 \gg PGE_2 \sim 12\text{-HHTrE} > PGD_2$ . Leukotrienes also regulate the inflammatory response in the airway/lung, notably during ARDS,<sup>17,18</sup> and their involvement in COVID-19 has been



**FIGURE 1** Levels of eicosanoids in the BAL fluids of healthy and COVID-19 patients. BALs were obtained and processed as described in *Methods*. Lipids then were extracted and quantitated by LC-MS/MS. A, Metabolism of arachidonic acid (AA) by the cyclooxygenase (COX) pathway. B, AA levels. C, D, Levels of COX-derived metabolites. E, Leukotriene (LT) biosynthetic pathways. F, G, Levels of LTs and exins (EX). H, Metabolism of AA and eicosapentaenoic acid (EPA) by 5-, 12-, and 15-lipoxygenase (LO). I, J, Levels of LO-derived metabolites of AA. K, EPA levels. L, Levels of LO-derived EPA metabolites. L, Proportions of eicosanoids from panel C, D, F, G, I, and K. Results are from the BALs of 25 healthy subjects and 33 COVID-19 patients. P values were obtained by performing a Mann-Whitney test using the GraphPad Prism 9 software: \*\*\*\* =  $P < .001$ , \*\*\*\*\* =  $P < .0001$




postulated.<sup>19</sup> Leukotrienes were not detected in the BALs of healthy volunteers, with the exception of LTB<sub>4</sub> (4/25 subjects). In contrast, COVID-19 patients had increased levels of LTB<sub>4</sub> and its CYP4F3A metabolite 20-COOH-LTB<sub>4</sub> (Figure 1F) as well as minor metabolites (20-OH-LTB<sub>4</sub> and 12-oxo-LTB<sub>4</sub>). Noteworthy, LTB<sub>4</sub> levels in the BALs of COVID-19 patients are comparable to those found in BALs of ARDS patients.<sup>20</sup> Moreover, the ratio between the sum of 20-OH- and 20-COOH-LTB<sub>4</sub> to LTB<sub>4</sub> ((20-OH- + 20-COOH-LTB<sub>4</sub>)/LTB<sub>4</sub>) was  $2.843 \pm 0.532$  (mean  $\pm$  SEM), reflecting an increased omega oxidation of LTB<sub>4</sub>. The cysteinyl-leukotrienes LTE<sub>4</sub> and the 15-LO-derived eoxin E<sub>4</sub> (EXE<sub>4</sub>) were also found in most COVID-19 patients (Figure 1G). Surprisingly, EXE<sub>4</sub> levels trended to be higher than those of LTE<sub>4</sub>. In line with the increase in both 5-LO- and 15-LO-derived leukotrienes, other 5-, 12-, and 15-LO metabolites, were increased in the BALs of COVID-19 patients, notably 5-HETE, 12-HETE, 15-HETE, and 15-KETE (Figure 1I,J). Dual LO metabolites from AA were also significantly increased. Of note, 12- and 15-HEPE derived from EPA metabolism, were also increased in the BAL of COVID-19 patients despite comparable levels of EPA between healthy controls and COVID-19 patients (Figure 1K,L), while the EPA-derived SPM RvE<sub>1</sub> was not detected. Overall, the eicosanoid profile of intubated


COVID-19 patients mainly consisted of TXB<sub>2</sub> and prostaglandins (62%), followed by the different HETEs and LTs (Figure 1L).

Many SPMs arise from the metabolism of docosanoids<sup>9</sup> and given their documented pro-resolving actions and antimicrobial activities, SPMs were recently postulated as being a potential approach for diminishing the inflammatory burden of COVID-19.<sup>21</sup> Both DPA(n-3) and DHA levels were increased in the BALs of COVID-19 patients (Figure 2B). We also observed significant increases in 17-HDPA(n-3), 14-HDHA and 17-HDHA (Figure 2C). In line with these findings, D-series resolvins (aside from RvD3) and Protectin D1 (PDX) were also increased in the BALs of COVID-19 patients (Figure 2D). Despite our good sensitivity (Table S1), Maresin-1 and -2 were not detected.

Lastly, we quantitated the levels of 15-LO metabolites derived from linoleic acid (LA), linolenic acid (ALA) and dihomo- $\gamma$ -linolenic acid (DGLA). LA levels were increased in the BAL of COVID-19 patients (Figure 3B), as well as its 15-LO-derived metabolites 13-HODE and 13-KODE (Figure 3C,D). Furthermore, 13-HOTrE (derived from ALA) and 15-HETrE (derived from DGLA) were both increased in COVID-19 patients (Figure 3D3C). We also observed an increase in the non-enzymatic metabolite 9-HODE (derived



**FIGURE 2** Levels of docosanoids in the BAL fluids of healthy and COVID-19 patients. BALs were obtained and processed as described in *Methods*. Lipids then were extracted and quantitated by LC-MS/MS. A, Docosanoid biosynthetic pathways. B, Levels of docosapentaenoic acid n-3 (DPA n-3) and docosahexaenoic acid (DHA). C,D, Docosanoid levels. E, Proportions of the different docosanoids from panel C and D. Results are from the BALs from 25 healthy subjects and 33 COVID-19 patients. P values were obtained by performing a Mann-Whitney test using the GraphPad Prism 9 software: \* =  $P < .05$ , \*\* =  $P < .01$ , \*\*\* =  $P < .001$ , \*\*\*\* =  $P < .0001$



**FIGURE 3** Levels of lipid mediators derived from linoleic acid (LA),  $\alpha$ -linolenic acid or dihomo- $\gamma$ -linolenic acid (DGLA) in the BAL fluids of healthy and COVID-19 patients. BALs were obtained and processed as described in *Methods*. Lipids then were extracted and quantitated by LC-MS/MS. A, Metabolism of LA, ALA, and DGLA by the 12/15-lipoxygenase (LO) pathways. B, Levels of LA. C,D, Levels of oxidized mediators from LA, ALA, and DGLA. Results are from the BALs from 25 healthy subjects and 33 COVID-19 patients. P values were obtained by performing a Mann-Whitney test using the GraphPad Prism 9 software: \*\*\*\* =  $P < .0001$

from LA) (Figure 3C). We also found that other auto-oxidative (or minor enzymatic) products were increased in the BAL of COVID-19 patients, namely 8-HETE and 11-HETE (both from AA) as well as 18-HEPE (from EPA), suggesting to an increased oxidative burst in the lungs of intubated COVID-19 patients (not shown).


Given that the increased in most lipids mediators we quantitated, we next addressed if correlations between lipids existed in the BAL of intubated COVID-19 patients. Most lipids consistently correlated with each other with few exceptions (Figure 4). Indeed, fatty acids levels were not a key determinant for the levels of their metabolites, indicating that downstream enzymes (cyclooxygenases, lipoxygenases) are likely more important in that respect. Interestingly, all fatty acids correlated with each other. As expected, lipids within the same metabolic pathway correlated with each other. Moreover, all 12- and 15-LO-derived mediators (HETEs, diHETEs, RvDs, and PDX) correlated with each other, regardless of their fatty acid of origin. Unsurprisingly, the levels of LXA<sub>4</sub> correlated with those of 5-HETE and 15-HETE, given that LXA<sub>4</sub> requires both the 5- and the 15-LO. Interestingly, LTE<sub>4</sub> and EXE<sub>4</sub> had a trend toward a negative correlation ( $r = -0.322$ ,  $P = .0676$ ) and they correlated with different bioactive lipid subsets, LTE<sub>4</sub> positively correlating with

fatty acid and 15-HETrE levels while EXE<sub>4</sub> mostly correlating with other 15-lipoxygenase metabolites and LTB<sub>4</sub>. This suggests that some COVID-19 patients are more prone to biosynthesize LTC<sub>4</sub> while others are more prone to biosynthesize EXC<sub>4</sub>.

We also investigated whether lipids from the BAL of COVID-19 patients correlated with the clinical parameters shown in Table 1 and found a very limited number of correlations (Figure 5). Fatty acids negatively correlated with AST, ALT and body weight (not significant for DPAn-3 and DHA), and positively correlated with the circulating platelet-to-lymphocyte ratio. LTE<sub>4</sub> negatively correlated with blood eosinophil counts and ALT, while positively correlating with hemoglobin. In contrast, EXE<sub>4</sub> positively correlated with blood eosinophils and we found a trend toward a positive correlation with ALT and toward an almost significant negative correlation with hemoglobin ( $r = -0.339$ ,  $P$ -value = 0.054). Interestingly, LXA<sub>4</sub>, PDX and, RvD4 positively correlated with blood eosinophil counts.

## 4 | DISCUSSION

Bioactive lipids are recognized pharmaceutical targets in numerous inflammatory diseases and lipid modulators have



**FIGURE 4** Correlation between bioactive lipid mediators in the BALs of COVID-19 patients. A Spearman correlation matrix between the different lipid mediators is shown. Positive correlations are shown using shades of blue.  $P < .05$  was the threshold of significance and non-significant correlations ( $P > .05$ ) are marked by an X


been utilized over the last fifty years with great success. Defining whether bioactive lipid levels were increased in severe COVID-19 patients was thus imperative. The lipidome arising from the metabolism of polyunsaturated fatty acids containing 20 carbons (eicosanoids) or 22 carbons (docosanoids) is continually expanding. While we performed a targeted lipidomic analysis on a significant fraction of them, we could not quantitate all of them, notably SPMs, due to our inability to secure appropriate internal standards. Thus, we cannot speculate on the levels and/or the potential roles of eicosanoids and docosanoids that we did not analyze in SARS-CoV-2-infected people.

Herein, we provide evidence that BALs of intubated COVID-19 patients contain extensive levels of bioactive lipids, which likely participate in the deleterious inflammatory response. More specifically, we highlight (1) an increase in COX metabolites, notably thromboxane; (2) an increase in leukotrienes, notably LTB<sub>4</sub>, LTE<sub>4</sub>, and EXE<sub>4</sub>; (3) an increase in 15-LO metabolites from LA, AA, EPA, DPA, and DHA;

(4) an increase in SPMs; (5) correlations between most lipid mediators, (6) a limited number of correlations between BAL lipids vs blood markers and/or clinical parameters.

The important increase in COX metabolites (Figure 1C) was expected, notably because of the previously documented increase in IL-1 $\beta$ ,<sup>20</sup> the latter being a good COX-2 inducer.<sup>22</sup> TXA<sub>2</sub> promotes bronchoconstriction and activate leukocytes and platelets, which is consistent with the enhanced platelet activation and thrombosis in COVID-19 and the strong link between TXB<sub>2</sub> and ARDS.<sup>10,23-26</sup> PGD<sub>2</sub> acting via the DP<sub>2</sub> receptor promotes the recruitment/activation of eosinophils, basophils, mastocytes, and innate lymphoid cells.<sup>27,28</sup> In contrast, PGE<sub>2</sub> is mostly anti-inflammatory via the EP<sub>2</sub> and EP<sub>4</sub> receptors<sup>29</sup> and PGI<sub>2</sub> is a recognized microvasculature dilator. In that respect, a retrospective observational study showed that the administration of PGI<sub>2</sub> (epoprostenol) was effective at improving the PaO<sub>2</sub>/FiO<sub>2</sub> ratio in 50% of COVID-19 patients.<sup>30</sup> Therapies targeting the prostaglandin pathway are numerous and usually effective at controlling





**FIGURE 5** Correlation between bioactive lipids in the BALs of COVID-19 patients and clinical parameters. A non-parametric Spearman correlation matrix between Lipid mediators and clinical parameters was done using the *corrplot* package. Positive correlations are shown using shades of blue and negative correlations are shown using shades of red.  $P < .05$  was the threshold of significance and non-significant correlations ( $P > .05$ ) are marked by an X

inflammation. Dexamethasone limits COX-2 expression<sup>22,31</sup> and might diminish COVID-19 severity and mortality, at least in part, by diminishing COX metabolites.<sup>32</sup> Aspirin was shown to significantly diminish ICU admissions and the need of mechanical ventilation,<sup>33</sup> and to diminish disease duration and mortality,<sup>34</sup> which is in line with a possible clinical benefit of early aspirin for the prevention of ARDS.<sup>26</sup> Altogether, our data support the concept that dexamethasone might improve severe COVID-19 by diminishing the prostaglandins/thromboxane storm in the lungs and that aspirin is also probably safe and beneficial for patients severely affected by COVID-19 although this needs to be investigated. To that end, aspirin would be a cheap and easily accessible alternative to limit the lipid storm and its putative burden during severe COVID-19 episodes. Finally, testing whether blocking the deleterious effects of PGD<sub>2</sub> and TXA<sub>2</sub> with the dual DP<sub>2</sub>/TP antagonist Ramatroban might be beneficial in COVID-19 is of interest, as proposed recently.<sup>35</sup>

Leukotrienes (LTB<sub>4</sub>, LTE<sub>4</sub>, and EXE<sub>4</sub>) were also increased in the BALs of COVID-19 patients (Figure 1F,G). This indicates that LTB<sub>4</sub> and cysteinyl-leukotrienes might also participate in the inflammatory burden of severe COVID-19 patients.<sup>19</sup> While LTB<sub>4</sub> is known to stimulate host defense during viral infections,<sup>35</sup> the observed  $\omega$ -LTB<sub>4</sub>/LTB<sub>4</sub> ratio of 2.84 is comparable to those found in cystic fibrosis and severely burned patients, two conditions in which infections

are poorly controlled.<sup>36,37</sup> Conceptually, this might diminish LTB<sub>4</sub> effectiveness at promoting its host defense-related functions.<sup>38</sup> The trend toward an inverse correlation between LTE<sub>4</sub> and EXE<sub>4</sub>, which is supported by their distinct correlation pattern with other lipids (Figure 4) and clinical parameters (Figure 5), suggests that these cysteinyl-LTs are coming from different cellular sources. To that end, EXE<sub>4</sub> levels in the BALs positively and significantly correlated with blood eosinophils counts while there was a strong trend between EXE<sub>4</sub> and BAL eosinophils counts. This suggests that eosin presence/biosynthesis is partly linked to eosinophils although the involvement of other 15-LO expressing cells, notably epithelial cells cannot be excluded.<sup>39</sup> Furthermore, the detection of EXE<sub>4</sub> in the BAL is of importance, since this mediator has rarely been detected in biological samples. Our data also suggest that the use of phosphodiesterase 4 inhibitors (eg, Roflumilast) in COVID-19 could be of interest as they might diminish inflammation by increasing intracellular cAMP concentrations, thereby decreasing AA release and eicosanoid biosynthesis.<sup>40-44</sup>

The metabolism of the omega-3 fatty acids EPA, DPA, and DHA notably leads to SPM biosynthesis, which are anti-inflammatory and pro-resolving lipids. Their high levels and correlations with eicosanoids were unexpected as they generally appear during the resolution of inflammation, after prostaglandin/leukotriene biosynthesis occurred, according to the



class-switch model.<sup>45</sup> However, our data indicate that prostaglandins, leukotrienes, and SPMs co-exist and participate in the inflammatory cascade during the acute phase of inflammation/infection in which resolution has not fully been engaged. Moreover, given their documented anti-inflammatory effects and their host-defense boosting functions, they could be viewed as potential mediators to enhance to limit the infection/inflammation. To that end, it was proposed that dexamethasone might promote some of its beneficial effects by upregulating SPM biosynthesis/effects in moderate to severe COVID-19 patients.<sup>46</sup> As of today, no resolution pharmacology treatment has been approved by governmental health agencies with the exception of EPA-ethyl ester (Vascepa). Nonetheless, EPA, DHA as well as some SPM precursors (18-HEPE, 17-HDHA) are available as dietary supplements. Finally, we must point out that only a fraction of docosanoids belonging to SPMs was analyzed and that other SPMs might also play a role during SARS-CoV-2 infections.

While it is largely accepted that the cytosolic phospholipase A<sub>2</sub> releases most of AA in activated human leukocytes (eg, in 47), the increased levels of LA, DPA, and DHA indicate that other phospholipases are also involved in the lipiome we found. The correlation between all fatty acids might also indicate that some fatty acids are released by the same phospholipases. Those phospholipases are likely secreted phospholipases A<sub>2</sub>, which can release fatty acids other than AA<sup>48,49</sup> and are contributing to lung inflammation<sup>50,51</sup> but our findings do not allow to pinpoint which additional phospholipases are involved.

One of the striking finding of this study is the very poor correlations between the lipid mediators from the BALs and the blood markers available. The significant negative correlations between liver enzymes and BAL fatty acids are puzzling and difficult to interpret as it is counterintuitive. While a liver-lung axis may promote damages in ARDS,<sup>52</sup> it is unclear to which extent such an axis participates in the pathogenesis and lipid mediator biosynthesis/effects in COVID-19.<sup>53,54</sup> Finally, even if blood TXB<sub>2</sub> levels are increased in COVID-19 patients,<sup>10</sup> those found in the plasma of our intubated COVID-19 patients did not correlate with those found in their BALs (data not shown), supporting the idea that the lipid mediator profile found in the lungs is not reflected in the circulation.

Our study has, however, some limitations. (1) BALs from healthy controls are from younger individuals than those from severe COVID-19 patients. This could have an impact on some mediators, notably PGD<sub>2</sub>, which participate in the worsening of infections mediated by SARS-CoV and are increased in aged mice<sup>55</sup>; (2) The procedure to obtain BAL fluids is slightly different between healthy controls and severe COVID-19 patients. Indeed, while lipid mediator analyses were done on the BAL return from one 50 mL saline bolus in healthy people, that of severe COVID-19 patients

were done on two boli. Although the possible outcome of this discrepancy could be and underestimation of lipid levels in one group as opposed to the other, these comparisons have permitted to estimate whether candidate lipids were to be expected in BAL in normal conditions. While some might argue that the first saline bolus contains more lipids than the following one(s), others might argue that lipid levels are consistent in each bolus. Given that, in our hands, the number of BAL cells is usually greater in the first bolus than in the subsequent ones (when multiple boli are utilized), we are confident that the BAL fluid from the first bolus likely contains more lipids than the second one. This would thus lead to a slight underestimation of lipids in the COVID-19 group. (3) Unfortunately, BAL fluid leukocyte counts of severe COVID-19 patients did not include the number of alveolar macrophages and it is thus impossible to determine a clear differential count of leukocytes in those samples. Furthermore, the eosinophil counts were somewhat limited by the apparatus as it could estimate the counts to  $\pm 10^5$ /mL cells, which likely impacts the different correlations involving eosinophils, notably with EXE<sub>4</sub>; (4) We could not compare the levels of all lipids in the BALs of severe COVID-19 patients to those found in the blood. This appears important given the limited number of correlations found between BAL lipids and most recognized COVID-19-related biomarkers such as CRP and D-dimers. To that end, a recent study indicated differences in some of the herein investigated lipids in the circulation.<sup>11</sup> It will thus be important to pursue our investigations and determine whether there is a transposition of the BAL lipiome to the circulation although our preliminary data indicate this is not the case.

In conclusion, our data indicate that lung fluids of severe COVID-19 patients, in addition to the cytokine storm previously documented, a lipid mediator storm also occurs. The correlations of most lipid mediators with each other regardless of their origin or biosynthetic pathways suggests an increase production of lipid mediators in general in the lungs of COVID-19 patients, as we observe for cytokine production. The rich library of clinically approved and low-cost lipid modulators might provide beneficial and important add-ons to the current therapeutic arsenal utilized to diminish COVID-19 severity and possibly that of other coronavirus-mediated infections.

## ACKNOWLEDGMENTS

ASA, AC, VR, ML, and NF are members of the Quebec Respiratory Health research Network. This work was supported by the Cheikh Zaid Foundation awarded to YZ and grants from the Canadian New Frontier Research Fund (NFRN-2019-00004) awarded to LF (PI), EB (co-PI) and NF (co-PI) and the Natural Sciences and Engineering Research Council of Canada awarded to NF. ASA received a doctoral award from the Canadian Institutes of Health Research. ED

and EB are respectfully recipient of doctoral and senior awards from the Fonds de Recherche du Québec en Santé (FRQS).


### CONFLICT OF INTEREST

The authors have stated explicitly that there are no conflicts of interest in connection with this article.

### AUTHOR CONTRIBUTIONS

ASA, YZ, YA, AC, HF, AEH, YT, AC, ML, EB, LF, and NF designed the research. ASA, YZ, VR, CT, ED, ID, CM, performed the research. ASA, YZ, OF, LF, and NF analyzed the data. All authors wrote the paper.

### ORCID

Anne-Sophie Archambault  <https://orcid.org/0000-0001-7243-7445>

Younes Zaid  <https://orcid.org/0000-0001-8750-9106>

Éric Boilard  <https://orcid.org/0000-0001-6319-6432>

Louis Flamand  <https://orcid.org/0000-0001-5010-4586>

Nicolas Flamand  <https://orcid.org/0000-0001-5944-7200>

### REFERENCES

1. Wu F, Zhao S, Yu B, et al. A new coronavirus associated with human respiratory disease in China. *Nature*. 2020;579:265-269.
2. Zhang XS, Duchaine C. SARS-CoV-2 and health care worker protection in low-risk settings: a review of modes of transmission and a novel airborne model involving inhalable particles. *Clin Microbiol Rev*. 2020;34.
3. Verdoni L, Mazza A, Gervasoni A, et al. An outbreak of severe Kawasaki-like disease at the Italian epicentre of the SARS-CoV-2 epidemic: an observational cohort study. *Lancet*. 2020;395:1771-1778.
4. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497-506.
5. Damiani S, Fiorentino M, De Palma A, et al. Pathological post mortem findings in lungs infected with Sars-Cov 2. *J Pathol*. 2020.
6. Barton LM, Duval EJ, Stroberg E, Ghosh S, Mukhopadhyay S. COVID-19 autopsies, Oklahoma, USA. *Am J Clin Pathol*. 2020;153:725-733.
7. Amigues I, Pearlman AH, Patel A, et al. Coronavirus disease 2019: investigational therapies in the prevention and treatment of hyperinflammation. *Expert Rev Clin Immunol*. 2020;16(12):1185-1204. <https://doi.org/10.1080/1744666X.2021.1847084>
8. Rubin EJ, Longo DL, Baden LR. Interleukin-6 receptor inhibition in Covid-19 - cooling the inflammatory soup. *N Engl J Med*. 2021;384:1564-1565.
9. Serhan CN, Levy BD. Resolvins in inflammation: emergence of the pro-resolving superfamily of mediators. *J Clin Invest*. 2018;128:2657-2669.
10. Zaid Y, Guessous F, Puhm F, et al. Platelet reactivity to thrombin differs between patients with COVID-19 and those with ARDS unrelated to COVID-19. *Blood Adv*. 2021;5:635-639.
11. Schwarz B, Sharma L, Roberts L, et al. Cutting edge: Severe SARS-CoV-2 infection in humans is defined by a shift in the serum lipidome, resulting in dysregulation of eicosanoid immune mediators. *J Immunol*. 2020.
12. Turcotte C, Dumais E, Archambault AS, et al. Human leukocytes differentially express endocannabinoid-glycerol lipases and hydrolyze 2-arachidonoyl-glycerol and its metabolites from the 15-lipoxygenase and cyclooxygenase pathways. *J Leukoc Biol*. 2019;106(6):1337-1347. <https://doi.org/10.1002/JLB.3A0919-049RRR>
13. Del Valle DM, Kim-Schulze S, Huang HH, et al. An inflammatory cytokine signature predicts COVID-19 severity and survival. *Nat Med*. 2020;26(10):1636-1643. <https://doi.org/10.1038/s41591-020-1051-9>
14. Tang Y, Liu J, Zhang D, Xu Z, Ji J, Wen C. Cytokine storm in COVID-19: The current evidence and treatment strategies. *Front Immunol*. 2020;11:1708.
15. Xiong Y, Liu Y, Cao L, et al. Transcriptomic characteristics of bronchoalveolar lavage fluid and peripheral blood mononuclear cells in COVID-19 patients. *Emerg Microbes Infect*. 2020;9:761-770.
16. Hu B, Huang S, Yin L. The cytokine storm and COVID-19. *J Med Virol*. 2021;93:250-256.
17. Stephenson AH, Lonigro AJ, Hyers TM, Webster RO, Fowler AA. Increased concentrations of leukotrienes in bronchoalveolar lavage fluid of patients with ARDS or at risk for ARDS. *Am Rev Respir Dis*. 1988;138:714-719.
18. Ratnoff WD, Matthay MA, Wong MY, et al. Sulfidopeptide-leukotriene peptidases in pulmonary edema fluid from patients with the adult respiratory distress syndrome. *J Clin Immunol*. 1988;8:250-258.
19. Funk CD, Ardakani A. A novel strategy to mitigate the hyperinflammatory response to COVID-19 by targeting leukotrienes. *Front Pharmacol*. 2020;11:1214.
20. Liao M, Liu Y, Yuan J, et al. Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19. *Nat Med*. 2020;26:842-844.
21. Panigrahy D, Gilligan MM, Huang S, et al. Inflammation resolution: a dual-pronged approach to averting cytokine storms in COVID-19? *Cancer Metastasis Rev*. 2020;39:337-340.
22. Mitchell JA, Belvisi MG, Akaraseenont P, et al. Induction of cyclo-oxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasone. *Br J Pharmacol*. 1994;113:1008-1014.
23. Zaid Y, Puhm F, Allaey I, et al. Platelets can associate with SARS-Cov-2 RNA and are hyperactivated in COVID-19. *Circ Res*. 2020;127:1404-1418.
24. Manne BK, Denorme F, Middleton EA, et al. Platelet gene expression and function in patients with COVID-19. *Blood*. 2020;136:1317-1329.
25. Hottz ED, Azevedo-Quintanilha IG, Palhinha L, et al. Platelet activation and platelet-monocyte aggregate formation trigger tissue factor expression in patients with severe COVID-19. *Blood*. 2020;136:1330-1341.
26. Abdulnour RE, Gunderson T, Barkas I, et al. Early intravascular events are associated with development of acute respiratory distress syndrome. a substudy of the LIPS-A clinical trial. *Am J Respir Crit Care Med*. 2018;197:1575-1585.
27. Hirai H, Tanaka K, Yoshie O, et al. Prostaglandin D2 selectively induces chemotaxis in T helper type 2 cells, eosinophils, and basophils via seven-transmembrane receptor CRTH2. *J Exp Med*. 2001;193:255-261.

28. Chang JE, Doherty TA, Baum R, Broide D. Prostaglandin D2 regulates human type 2 innate lymphoid cell chemotaxis. *J Allergy Clin Immunol.* 2014;133(899–901):e893.
29. Medeiros A, Peres-Buzalaf C, Fortino Verdan F, Serezani CH. Prostaglandin E<sub>2</sub> and the suppression of phagocyte innate immune responses in different organs. *Mediators Inflamm.* 2012;2012:327568.
30. Sonti R, Pike CW, Cobb N. Responsiveness of inhaled epoprostenol in respiratory failure due to COVID-19. *J Intensive Care Med.* 2020;885066620976525.
31. Crofford LJ, Wilder RL, Ristimaki AP, et al. Cyclooxygenase-1 and -2 expression in rheumatoid synovial tissues. Effects of interleukin-1b, phorbol ester, and corticosteroids. *J Clin Invest.* 1994;93:1095-1101.
32. Tomazini BM, Maia IS, Cavalcanti AB, et al. Effect of dexamethasone on days alive and ventilator-free in patients with moderate or severe acute respiratory distress syndrome and COVID-19: the CoDEX randomized clinical trial. *JAMA.* 2020;324:1307-1316.
33. Chow JH, Khanna AK, Kethireddy S, et al. Aspirin use is associated with decreased mechanical ventilation, ICU admission, and in-hospital mortality in hospitalized patients with COVID-19. *Anesth Analg.* 2020;132(4):930-941.
34. Merzon E, Green I, Vinker S, et al. The use of aspirin for primary prevention of cardiovascular disease is associated with a lower likelihood of COVID-19 infection. *FEBS J.* 2021. <https://doi.org/10.1111/febs.15784>
35. Gupta A, Kalantar-Zadeh K, Reddy ST. Ramatroban as a novel immunotherapy for COVID-19. *J Mol Genet Med.* 2020;14.
36. Yang J, Eiserich JP, Cross CE, Morrissey BM, Hammock BD. Metabolomic profiling of regulatory lipid mediators in sputum from adult cystic fibrosis patients. *Free Radic Biol Med.* 2012;53:160-171.
37. Brom J, Konig W, Koller M, Gross-Weege W, Erbs G, Muller F. Metabolism of leukotriene B<sub>4</sub> by polymorphonuclear granulocytes of severely burned patients. *Prostaglandins, Leukot Med.* 1987;27:209-225.
38. Archambault AS, Poirier S, Lefebvre JS, et al. 20-Hydroxy- and 20-carboxy-leukotriene (LT) B<sub>4</sub> downregulate LTB<sub>4</sub>-mediated responses of human neutrophils and eosinophils. *J Leukoc Biol.* 2019;105:1131-1142.
39. Brunnstrom A, Tryselius Y, Feltenmark S, et al. On the biosynthesis of 15-HETE and eoxin C<sub>4</sub> by human airway epithelial cells. *Prostaglandins Other Lipid Mediat.* 2015;121:83-90.
40. Ham EA, Soderman DD, Zanetti ME, Dougherty HW, McCauley E, Kuehl FA Jr. Inhibition by prostaglandins of leukotriene B<sub>4</sub> release from activated neutrophils. *Proc Natl Acad Sci USA.* 1983;80:4349-4353.
41. Fonteh AN, Winkler JD, Torphy TJ, Heravi J, Udem BJ, Chilton FH. Influence of isoproterenol and phosphodiesterase inhibitors on platelet-activating factor biosynthesis in the human neutrophil. *J Immunol.* 1993;151:339-350.
42. Denis D, Riendeau D. Phosphodiesterase 4-dependent regulation of cyclic AMP levels and leukotriene B<sub>4</sub> biosynthesis in human polymorphonuclear leukocytes. *Eur J Pharmacol.* 1999;367:343-350.
43. Flamand N, Boudreault S, Picard S, et al. Adenosine, a potent natural suppressor of arachidonic acid release and leukotriene biosynthesis in human neutrophils. *Am J Respir Crit Care Med.* 2000;161:S88-94.
44. Flamand N, Plante H, Picard S, Laviolette M, Borgeat P. Histamine-induced inhibition of leukotriene biosynthesis in human neutrophils: involvement of the H<sub>2</sub> receptor and cAMP. *Br J Pharmacol.* 2004;141:552-561.
45. Buckley CD, Gilroy DW, Serhan CN. Proresolving lipid mediators and mechanisms in the resolution of acute inflammation. *Immunity.* 2014;40:315-327.
46. Andreakos E, Papadaki M, Serhan CN. Dexamethasone, proresolving lipid mediators and resolution of inflammation in COVID-19. *Allergy.* 2020.
47. Flamand N, Picard S, Lemieux L, Pouliot M, Bourgoin SG, Borgeat P. Effects of pyrrophenone, an inhibitor of group IVA phospholipase A<sub>2</sub>, on eicosanoid and PAF biosynthesis in human neutrophils. *Br J Pharmacol.* 2006;149:385-392.
48. Hanasaki K, Ono T, Saiga A, et al. Purified group X secretory phospholipase A<sub>2</sub> induced prominent release of arachidonic acid from human myeloid leukemia cells. *J Biol Chem.* 1999;274:34203-34211.
49. Murakami M, Masuda S, Shimbara S, et al. Cellular arachidonate-releasing function of novel classes of secretory phospholipase A<sub>2</sub>s (groups III and XII). *J Biol Chem.* 2003;278:10657-10667.
50. Hallstrand TS, Lai Y, Altemeier WA, et al. Regulation and function of epithelial secreted phospholipase A<sub>2</sub> group X in asthma. *Am J Respir Crit Care Med.* 2013;188:42-50.
51. Arbibe L, Koumanov K, Vial D, et al. Generation of lysophospholipids from surfactant in acute lung injury is mediated by type-II phospholipase A<sub>2</sub> and inhibited by a direct surfactant protein A-phospholipase A<sub>2</sub> protein interaction. *J Clin Invest.* 1998;102:1152-1160.
52. Herrero R, Sanchez G, Asensio I, et al. Liver-lung interactions in acute respiratory distress syndrome. *Intensive Care Med Exp.* 2020;8:48.
53. Weber S, Hellmuth JC, Scherer C, Muenchhoff M, Mayerle J, Gerbes AL. Liver function test abnormalities at hospital admission are associated with severe course of SARS-CoV-2 infection: a prospective cohort study. *Gut.* 2021.
54. Portincasa P, Krawczyk M, Machill A, Lammert F, Di Ciaula A. Hepatic consequences of COVID-19 infection. Lapping or biting? *Eur J Intern Med.* 2020;77:18-24.
55. Zhao J, Zhao J, Legge K, Perlman S. Age-related increases in PGD<sub>2</sub> expression impair respiratory DC migration, resulting in diminished T cell responses upon respiratory virus infection in mice. *J Clin Invest.* 2011;121:4921-4930.

## SUPPORTING INFORMATION

Additional Supporting Information may be found online in the Supporting Information section.

**How to cite this article:** Archambault A-S, Zaid Y, Rakotoarivelo V, et al. High levels of eicosanoids and docosanoids in the lungs of intubated COVID-19 patients. *The FASEB Journal.* 2021;35:e21666. <https://doi.org/10.1096/fj.202100540R>